

## Geometric Progression

---

### Question 1.

Find, which of the following sequence form a G.P. :

(i) 8, 24, 72, 216, .....

(ii)  $\frac{1}{8}, \frac{1}{24}, \frac{1}{72}, \frac{1}{216}, \dots\dots\dots$

(iii) 9, 12, 16, 24, .....

### Solution 1(i).

Given sequence: 8, 24, 72, 216.....

Now,

$$\frac{24}{8} = 3, \quad \frac{72}{24} = 3, \quad \frac{216}{72} = 3$$

Since  $\frac{24}{8} = \frac{72}{24} = \frac{216}{72} = \dots\dots = 3$ , the given sequence is a G.P.

with common ratio 3.

### Solution 1(ii).

Given sequence:  $\frac{1}{8}, \frac{1}{24}, \frac{1}{72}, \frac{1}{216}, \dots\dots\dots$

Now,

$$\frac{\frac{1}{24}}{\frac{1}{8}} = \frac{1}{3}, \quad \frac{\frac{1}{72}}{\frac{1}{24}} = \frac{1}{3}, \quad \frac{\frac{1}{216}}{\frac{1}{72}} = \frac{1}{3}$$

Since  $\frac{\frac{1}{24}}{\frac{1}{8}} = \frac{\frac{1}{72}}{\frac{1}{24}} = \frac{\frac{1}{216}}{\frac{1}{72}} = \dots\dots = \frac{1}{3}$ , the given sequence is a G.P.

with common ratio  $\frac{1}{3}$ .

### Solution 1(iii).

Given sequence: 9, 12, 16, 24.....

Now,

$$\frac{12}{9} = \frac{4}{3}, \quad \frac{16}{12} = \frac{4}{3}, \quad \frac{24}{16} = \frac{3}{2}$$

Since  $\frac{24}{9} = \frac{72}{24} \neq \frac{216}{72}$ , the given sequence is not a G.P.

**Question 2.**

Find the 9th term of the series :

1, 4, 16, 64 .....

**Solution:**

Given sequence: 1, 4, 16, 64.....

Now,

$$\frac{4}{1} = 4, \quad \frac{16}{4} = 4, \quad \frac{64}{16} = 4$$

Since  $\frac{4}{1} = \frac{16}{4} = \frac{64}{16} = \dots = 4$ , the given sequence is a G.P.

with first term,  $a = 1$  and common ratio,  $r = 4$ .

Now,  $t_n = ar^{n-1}$

$$\Rightarrow t_9 = 1 \times 4^8 = 65536$$

**Question 3.**

Find the seventh term of the G.P. :

1,  $\sqrt{3}$ , 3,  $3\sqrt{3}$  .....

**Solution:**

Given G.P.: 1,  $\sqrt{3}$ , 3,  $3\sqrt{3}$ , .....

Here,

First term,  $a = 1$

Common ratio,  $r = \frac{\sqrt{3}}{1} = \sqrt{3}$

Now,  $t_n = ar^{n-1}$

$$\Rightarrow t_7 = 1 \times (\sqrt{3})^6 = 27$$

**Question 4.**

Find the 8<sup>th</sup> term of the sequence :

$\frac{3}{4}$ ,  $1\frac{1}{2}$ , 3, .....

**Solution:**

Given sequence:  $\frac{3}{4}, 1\frac{1}{2}, 3, \dots$

i.e.  $\frac{3}{4}, \frac{3}{2}, 3, \dots$

Now,

$$\frac{\frac{3}{2}}{\frac{3}{4}} = 2, \quad \frac{3}{\frac{3}{2}} = 2,$$

Since  $\frac{\frac{3}{2}}{\frac{3}{4}} = \frac{3}{\frac{3}{2}} = \dots = 2$ , the given sequence is a G.P.

with first term,  $a = \frac{3}{4}$  and common ratio,  $r = 2$ .

Now,  $t_n = ar^{n-1}$

$$\Rightarrow t_8 = \frac{3}{4} \times 2^7 = \frac{3}{4} \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 \times 2 = 3 \times 2^5 = 96$$

**Question 5.**

Find the 10<sup>th</sup> term of the G.P. :

**Solution:**

Given G.P.:  $12, 4, 1\frac{1}{3}, \dots$

Here,

First term,  $a = 12$

Common ratio,  $r = \frac{4}{12} = \frac{1}{3}$

Now,  $t_n = ar^{n-1}$

$$\Rightarrow t_{10} = 12 \times \left(\frac{1}{3}\right)^9 = 12 \times \frac{1}{19683} = \frac{4}{6561}$$

**Question 6.**

Find the n<sup>th</sup> term of the series :

**Solution:**

Given series: 1, 2, 4, 8, .....

Now,

$$\frac{2}{1} = 2, \quad \frac{4}{2} = 2, \quad \frac{8}{4} = 2$$

Since  $\frac{2}{1} = \frac{4}{2} = \frac{8}{4} = \dots = 2$ , the given sequence is a G.P.

with first term,  $a = 1$  and common ratio,  $r = 2$ .

Now,  $t_n = ar^{n-1}$

$$\Rightarrow t_n = 1 \times 2^{n-1} = 2^{n-1}$$

**Question 7.**

Find the next three terms of the sequence :

$\sqrt{5}, 5, 5\sqrt{5}, \dots$

**Solution:**

Given sequence:  $\sqrt{5}, 5, 5\sqrt{5}, \dots$

Now,

$$\frac{5}{\sqrt{5}} = \sqrt{5}, \quad \frac{5\sqrt{5}}{5} = \sqrt{5}$$

Since  $\frac{5}{\sqrt{5}} = \frac{5\sqrt{5}}{5} = \dots = \sqrt{5}$ , the given sequence is a G.P.

with first term,  $a = \sqrt{5}$  and common ratio,  $r = \sqrt{5}$ .

Now,  $t_n = ar^{n-1}$

$\therefore$  Next three terms:

$$4^{\text{th}} \text{ term} = \sqrt{5} \times (\sqrt{5})^3 = \sqrt{5} \times 5\sqrt{5} = 25$$

$$5^{\text{th}} \text{ term} = \sqrt{5} \times (\sqrt{5})^4 = \sqrt{5} \times 25 = 25\sqrt{5}$$

$$6^{\text{th}} \text{ term} = \sqrt{5} \times (\sqrt{5})^5 = \sqrt{5} \times 25\sqrt{5} = 125$$

**Question 8.**

Find the sixth term of the series :

$2^2, 2^3, 2^4, \dots$

**Solution:**

Given sequence:  $2^2, 2^3, 2^4, \dots$

Now,

$$\frac{2^3}{2^2} = 2, \quad \frac{2^4}{2^3} = 2$$

Since  $\frac{2^3}{2^2} = \frac{2^4}{2^3} = \dots = 2$ , the given sequence is a G.P.

with first term,  $a = 2^2 = 4$  and common ratio,  $r = 2$ .

Now,  $t_n = ar^{n-1}$

$$\therefore t_6 = 4 \times (2)^5 = 4 \times 32 = 128$$

**Question 9.**

Find the seventh term of the G.P. :

$[\text{late}] \sqrt{3} + 1, 1, \frac{\sqrt{3} - 1}{2} [\text{late}]$ , .....

**Solution:**

Given G.P.:  $\sqrt{3} + 1, 1, \frac{\sqrt{3} - 1}{2}, \dots$

Here,

First term,  $a = \sqrt{3} + 1$

Common ratio,  $r = \frac{1}{\sqrt{3} + 1}$

Now,  $t_n = ar^{n-1}$

$$\Rightarrow t_7 = (\sqrt{3} + 1) \times \left( \frac{1}{\sqrt{3} + 1} \right)^6$$

$$= \left( \frac{1}{\sqrt{3} + 1} \right)^5$$

$$= \left( \frac{1}{\sqrt{3} + 1} \times \frac{\sqrt{3} - 1}{\sqrt{3} - 1} \right)^5$$

$$= \left( \frac{\sqrt{3} - 1}{2} \right)^5$$

$$= \frac{1}{32} (\sqrt{3} - 1)^5$$

**Question 10.**

Find the G.P. whose first term is 64 and next term is 32.

**Solution:**

First term,  $a = 64$

Second term,  $t_2 = 32$

$$\Rightarrow ar = 32$$

$$\Rightarrow 64 \times r = 32$$

$$\Rightarrow r = \frac{32}{64} = \frac{1}{2}$$

$\therefore$  Required G.P. =  $a, ar, ar^{n-1}, ar^{n-2}, \dots$

$$= 64, 32, 64 \times \left(\frac{1}{2}\right)^2, 64 \times \left(\frac{1}{2}\right)^3, \dots$$

$$= 64, 32, 16, 8, \dots$$

**Question 11.**

Find the next three terms of the series:

$$\frac{2}{27}, \frac{2}{9}, \frac{2}{3}, \dots$$

**Solution:**

Given sequence:  $\frac{2}{27}, \frac{2}{9}, \frac{2}{3}, \dots$

Now,

$$\frac{\frac{2}{9}}{\frac{2}{27}} = 3, \quad \frac{\frac{2}{3}}{\frac{2}{9}} = 3$$

Since  $\frac{\frac{2}{9}}{\frac{2}{27}} = \frac{\frac{2}{3}}{\frac{2}{9}} = \dots = 3$ , the given sequence is a G.P.

with first term,  $a = \frac{2}{27}$  and common ratio,  $r = 3$ .

Now,  $t_n = ar^{n-1}$

$\therefore$  Next three terms:

$$4^{\text{th}} \text{ term} = \frac{2}{27} \times (3)^3 = \frac{2}{27} \times 27 = 2$$

$$5^{\text{th}} \text{ term} = \frac{2}{27} \times (3)^4 = \frac{2}{27} \times 27 \times 3 = 6$$

$$6^{\text{th}} \text{ term} = \frac{2}{27} \times (3)^5 = \frac{2}{27} \times 27 \times 9 = 18$$

**Question 12.**

Find the next two terms of the series

$2 - 6 + 18 - 54 \dots\dots\dots$

**Solution:**

Given series:  $2 - 6 + 18 - 54 \dots\dots\dots$

Now,

$$\frac{-6}{2} = -3, \quad \frac{18}{-6} = -3, \quad \frac{-54}{18} = -3$$

Since  $\frac{-6}{2} = \frac{18}{-6} = \frac{-54}{18} = \dots\dots\dots = -3$ , the given sequence is a G.P.

with first term,  $a = 2$  and common ratio,  $r = -3$ .

Now,  $t_n = ar^{n-1}$

$\therefore$  Next two terms:

$$5^{\text{th}} \text{ term} = 2 \times (-3)^4 = 2 \times 81 = 162$$

$$6^{\text{th}} \text{ term} = 2 \times (-3)^5 = 2 \times (-243) = -486$$

**Exercise 11B****Question 1.**

Which term of the G.P. :

$$-10, \frac{5}{\sqrt{3}}, -\frac{5}{6}, \dots\dots\dots \text{ is } -\frac{5}{72} ?$$

**Solution:**

For the given G.P.:

First term,  $a = -10$

$$\text{Common ratio, } r = \frac{5/\sqrt{3}}{-10} = -\frac{1}{2\sqrt{3}}$$

If  $-\frac{5}{72}$  is the  $n^{\text{th}}$  term of the given G.P., then

$$-\frac{5}{72} = ar^{n-1}$$

$$\Rightarrow -\frac{5}{72} = -10 \times \left(\frac{1}{2\sqrt{3}}\right)^{n-1}$$

$$\Rightarrow \frac{1}{144} = \left(\frac{1}{2\sqrt{3}}\right)^{n-1}$$

$$\Rightarrow \frac{1}{2 \times 2 \times 2 \times 2 \times \sqrt{3} \times \sqrt{3} \times \sqrt{3} \times \sqrt{3}} = \left(\frac{1}{2\sqrt{3}}\right)^{n-1}$$

$$\Rightarrow \left(\frac{1}{2\sqrt{3}}\right)^4 = \left(\frac{1}{2\sqrt{3}}\right)^{n-1}$$

$$\Rightarrow n - 1 = 4$$

$$\Rightarrow n = 5$$

**Question 2.**

The fifth term of a G.P. is 81 and its second term is 24. Find the geometric progression.

**Solution:**

Let the first term of the G.P. be  $a$  and its common ratio be  $r$ .

$$5^{\text{th}} \text{ term} = 81 \Rightarrow ar^4 = 81$$

$$2^{\text{nd}} \text{ term} = 24 \Rightarrow ar = 24$$

$$\text{Now, } \frac{ar^4}{ar} = \frac{81}{24}$$

$$\Rightarrow r^3 = \frac{27}{8}$$

$$\Rightarrow r = \frac{3}{2}$$

$$ar = 24$$

$$\Rightarrow a = 16$$

$$\begin{aligned}\therefore \text{GP.} &= a, ar, ar^2, ar^3, \dots \\ &= 16, 24, 16 \times \left(\frac{3}{2}\right)^2, 16 \times \left(\frac{3}{2}\right)^3, \dots \\ &= 16, 24, 36, 54, \dots\end{aligned}$$

### Question 3.

Fourth and seventh terms of a G.P. are  $\frac{1}{18}$  and  $-\frac{1}{486}$  respectively. Find the GP.

#### Solution:

Let the first term of the G.P. be  $a$  and its common ratio be  $r$ .

$$4^{\text{th}} \text{ term} = \frac{1}{18} \Rightarrow ar^3 = \frac{1}{18}$$

$$7^{\text{th}} \text{ term} = -\frac{1}{486} \Rightarrow ar^6 = -\frac{1}{486}$$

$$\text{Now, } \frac{ar^6}{ar^3} = \frac{-\frac{1}{486}}{\frac{1}{18}}$$

$$\Rightarrow r^3 = -\frac{1}{27}$$

$$\Rightarrow r = -\frac{1}{3}$$

$$ar^3 = \frac{1}{18}$$

$$\Rightarrow a \times \left(-\frac{1}{3}\right)^3 = \frac{1}{18}$$

$$\Rightarrow a = -\frac{27}{18} = -\frac{3}{2}$$

$$\begin{aligned}\therefore \text{GP.} &= a, ar, ar^2, ar^3, \dots \\ &= -\frac{3}{2}, -\frac{3}{2} \times \left(\frac{-1}{3}\right), -\frac{3}{2} \times \left(\frac{-1}{3}\right)^2, \frac{1}{18}, \dots \\ &= -\frac{3}{2}, \frac{1}{2}, -\frac{1}{6}, \frac{1}{18}, \dots\end{aligned}$$

**Question 4.**

If the first and the third terms of a G.P. are 2 and 8 respectively, find its second term.

**Solution:**

Let the first term of the G.P. be  $a$  and its common ratio be  $r$ .

$$\therefore 1^{\text{st}} \text{ term} = a = 2$$

$$\text{And, } 3^{\text{rd}} \text{ term} = 8 \Rightarrow ar^2 = 8$$

$$\text{Now, } \frac{ar^2}{a} = \frac{8}{2}$$

$$\Rightarrow r^2 = 4$$

$$\Rightarrow r = \pm 2$$

When  $a = 2$  and  $r = 2$

$$2^{\text{nd}} \text{ term} = ar = 2 \times 2 = 4$$

When  $a = 2$  and  $r = -2$

$$2^{\text{nd}} \text{ term} = ar = 2 \times (-2) = -4$$

**Question 5.**

The product of 3rd and 8th terms of a G.P. is 243. If its 4th term is 3, find its 7th term.

**Solution:**

Let the first term of the G.P. be  $a$  and its common ratio be  $r$ .

Now,

$$t_3 \times t_8 = 243$$

$$\Rightarrow ar^2 \times ar^7 = 243$$

$$\Rightarrow a^2 r^9 = 243 \quad \dots(i)$$

Also,

$$t_4 = 3$$

$$\Rightarrow ar^3 = 3$$

$$\Rightarrow a = \frac{3}{r^3}$$

Substituting the value of  $a$  in (i), we get

$$\left(\frac{3}{r^3}\right)^2 \times r^9 = 243$$

$$\Rightarrow \frac{9}{r^6} \times r^9 = 243$$

$$\Rightarrow r^3 = 27$$

$$\Rightarrow r = 3$$

$$\Rightarrow a = \frac{3}{3^3} = \frac{3}{27} = \frac{1}{9}$$

$$\therefore 7^{\text{th}} \text{ term} = t_7 = ar^6 = \frac{1}{9} \times (3)^6 = 81$$

### Question 6.

Find the geometric progression with 4<sup>th</sup> term = 54 and 7<sup>th</sup> term = 1458.

### Solution:

Let the first term of the G.P. be  $a$  and its common ratio be  $r$ .

$$4^{\text{th}} \text{ term} = 54 \Rightarrow ar^3 = 54$$

$$7^{\text{th}} \text{ term} = 1458 \Rightarrow ar^6 = 1458$$

$$\text{Now, } \frac{ar^6}{ar^3} = \frac{1458}{54}$$

$$\Rightarrow r^3 = 27$$

$$\Rightarrow r = 3$$

$$ar^3 = 54$$

$$\Rightarrow a \times (3)^3 = 54$$

$$\Rightarrow a = \frac{54}{27} = 2$$

$$\begin{aligned} \therefore \text{G.P.} &= a, ar, ar^2, ar^3, \dots \\ &= 2, 2 \times 3, 2 \times (3)^2, 54, \dots \\ &= 2, 6, 18, 54, \dots \end{aligned}$$

### Question 7.

Second term of a geometric progression is 6 and its fifth term is 9 times of its third term. Find the geometric progression. Consider that each term of the G.P. is positive.

**Solution:**

Let the first term of the G.P. be  $a$  and its common ratio be  $r$ .

$$\text{Now, 2}^{\text{nd}} \text{ term} = t_2 = 6 \Rightarrow ar = 6$$

$$\text{Also, } t_5 = 9 \times t_3$$

$$\Rightarrow ar^4 = 9 \times ar^2$$

$$\Rightarrow r^2 = 9$$

$$\Rightarrow r = \pm 3$$

Since, each term of a G.P. is positive, we have  $r = 3$

$$ar = 6$$

$$\Rightarrow a \times 3 = 6 \Rightarrow a = 2$$

$$\begin{aligned} \therefore \text{G.P.} &= a, ar, ar^2, ar^3, \dots \\ &= 2, 6, 2 \times (3)^2, 2 \times (3)^3, \dots \\ &= 2, 6, 18, 54, \dots \end{aligned}$$

**Question 8.**

The fourth term, the seventh term and the last term of a geometric progression are 10, 80 and 2560 respectively. Find its first term, common ratio and number of terms.

**Solution:**

Let the first term of the G.P. be  $a$  and its common ratio be  $r$ .

Now,

$$4^{\text{th}} \text{ term} = t_4 = 10 \Rightarrow ar^3 = 10$$

$$7^{\text{th}} \text{ term} = t_7 = 80 \Rightarrow ar^6 = 80$$

$$\frac{ar^6}{ar^3} = \frac{80}{10}$$

$$\Rightarrow r^3 = 8$$

$$\Rightarrow r = 2$$

$$ar^3 = 10$$

$$\Rightarrow a \times (2)^3 = 10$$

$$\Rightarrow a = \frac{10}{8} = \frac{5}{4}$$

Last term =  $l = 2560$

Let there be  $n$  terms in given G.P.

$$\Rightarrow t_n = 2560$$

$$\Rightarrow ar^{n-1} = 2560$$

$$\Rightarrow \frac{5}{4} \times (2)^{n-1} = 2560$$

$$\Rightarrow (2)^{n-1} = 2048$$

$$\Rightarrow (2)^{n-1} = (2)^{11}$$

$$\Rightarrow n - 1 = 11$$

$$\Rightarrow n = 12$$

Thus, we have

First term =  $\frac{5}{4}$ , Common ratio = 2 and Number of terms = 12

### Question 9.

If the 4th and 9th terms of a G.P. are 54 and 13122 respectively, find the GP. Also, find its general term.

### Solution:

Let the first term of the G.P. be  $a$  and its common ratio be  $r$ .

Now,

$$4^{\text{th}} \text{ term} = t_4 = 54 \Rightarrow ar^3 = 54$$

$$9^{\text{th}} \text{ term} = t_9 = 13122 \Rightarrow ar^8 = 13122$$

$$\frac{ar^8}{ar^3} = \frac{13122}{54}$$

$$\Rightarrow r^5 = 243$$

$$\Rightarrow r = 3$$

$$ar^3 = 54$$

$$\Rightarrow a \times (3)^3 = 54$$

$$\Rightarrow a = \frac{54}{27} = 2$$

$$\begin{aligned} \therefore \text{Required G.P.} &= a, ar, ar^2, ar^3, \dots \\ &= 2, 2 \times 3, 2 \times (3)^2, 54 \\ &= 2, 6, 18, 54 \end{aligned}$$

$$\text{General term} = t_n = ar^{n-1} = 2 \times (3)^{n-1}$$

**Question 10.**

The fifth, eighth and eleventh terms of a geometric progression are p, q and r respectively. Show that :  $q^2 = pr$ .

**Solution:**

Let the first term of the G.P. be a and its common ratio be r.

$$5^{\text{th}} \text{ term} = t_5 = p$$

$$\Rightarrow ar^4 = p$$

$$8^{\text{th}} \text{ term} = t_8 = q$$

$$\Rightarrow ar^7 = q$$

$$11^{\text{th}} \text{ term} = t_{11} = r$$

$$\Rightarrow ar^{10} = r$$

Now,

$$pr = ar^4 \times ar^{10} = a^2 \times r^{14} = (a \times r^7)^2 = q^2$$

$$\Rightarrow q^2 = pr$$

**Exercise 11C****Question 1.**

Find the seventh term from the end of the series :  $\sqrt{2}, 2, 2\sqrt{2}, \dots, 32$ .

**Solution:**

Given series:  $\sqrt{2}, 2, 2\sqrt{2}, \dots, 32$

$$\text{Now, } \frac{2}{\sqrt{2}} = \sqrt{2}, \quad \frac{2\sqrt{2}}{2} = \sqrt{2}$$

So, the given series is a G.P. with common ratio,  $r = \sqrt{2}$

Here, last term,  $l = 32$

$$\therefore 7^{\text{th}} \text{ term from an end} = \frac{l}{r^6} = \frac{32}{(\sqrt{2})^6} = \frac{32}{8} = 4$$

**Question 2.**

Find the third term from the end of the GP.

$$\frac{2}{27}, \frac{2}{9}, \frac{2}{3}, \dots, 162$$

**Solution:**

Given G.P.:  $\frac{2}{27}, \frac{2}{9}, \frac{2}{3}, \dots, 162$

Here,

$$\text{Common ratio, } r = \frac{\frac{2}{9}}{\frac{2}{27}} = 3$$

Last term,  $l = 162$

$$\therefore 3^{\text{rd}} \text{ term from an end} = \frac{l}{r^2} = \frac{162}{(3)^2} = \frac{162}{9} = 18$$

**Question 3.**

For the  $\frac{1}{27}, \frac{1}{9}, \frac{1}{3}, \dots, 81$ ;

find the product of fourth term from the beginning and the fourth term from the end.

**Solution:**

Given G.P.:  $\frac{1}{27}, \frac{1}{9}, \frac{1}{3}, \dots, 81$

Here,

$$\text{Common ratio, } r = \frac{\frac{1}{9}}{\frac{1}{27}} = 3$$

First term,  $a = \frac{1}{27}$  and Last term,  $l = 81$

$$\therefore 4^{\text{th}} \text{ term from the beginning} = ar^3 = \frac{1}{27} \times (3)^3 = \frac{1}{27} \times 27 = 1$$

$$\text{And, } 4^{\text{th}} \text{ term from an end} = \frac{l}{r^3} = \frac{81}{(3)^3} = \frac{81}{27} = 3$$

Thus, required product =  $1 \times 3 = 3$

**Question 4.**

If for a G.P.,  $p^{\text{th}}$ ,  $q^{\text{th}}$  and  $r^{\text{th}}$  terms are  $a$ ,  $b$  and  $c$  respectively ; prove that :  
 $(q - r) \log a + (r - p) \log b + (p - q) \log c = 0$

**Solution:**

Let the first term of the G.P. be  $A$  and its common ratio be  $R$ .

Then,

$$p^{\text{th}} \text{ term} = a \Rightarrow AR^{p-1} = a$$

$$q^{\text{th}} \text{ term} = b \Rightarrow AR^{q-1} = b$$

$$r^{\text{th}} \text{ term} = c \Rightarrow AR^{r-1} = c$$

Now,

$$\begin{aligned} a^{q-r} \times b^{r-p} \times c^{p-q} &= (AR^{p-1})^{q-r} \times (AR^{q-1})^{r-p} \times (AR^{r-1})^{p-q} \\ &= A^{q-r} \cdot R^{(p-1)(q-r)} \times A^{r-p} \cdot R^{(q-1)(r-p)} \times A^{p-q} \cdot R^{(r-1)(p-q)} \\ &= A^{q-r+r-p+p-q} \times R^{(p-1)(q-r)+(q-1)(r-p)+(r-1)(p-q)} \\ &= A^0 \times R^0 \\ &= 1 \end{aligned}$$

Taking log on both the sides, we get

$$\log(a^{q-r} \times b^{r-p} \times c^{p-q}) = \log 1$$

$$\Rightarrow (q-r)\log a + (r-p)\log b + (p-q)\log c = 0 \quad \dots(\text{proved})$$

**Question 5.**

If  $a$ ,  $b$  and  $c$  in G.P., prove that :  $\log a^n$ ,  $\log b^n$  and  $\log c^n$  are in A.P.

**Solution:**

Here,  $a$ ,  $b$ ,  $c$  are in G.P.

$$\Rightarrow b^2 = ac$$

Taking log on both sides, we get

$$\log(b^2) = \log(ac)$$

$$\Rightarrow 2\log b = \log a + \log c$$

$$\Rightarrow \log b + \log b = \log a + \log c$$

$$\Rightarrow \log b - \log a = \log c - \log b$$

$$\Rightarrow \log a, \log b \text{ and } \log c \text{ are in A.P.}$$

**Question 6.**

If each term of a G.P. is raised to the power  $x$ , show that the resulting sequence is also a G.P.

**Solution:**

Let  $a_1, a_2, a_3, \dots, a_n, \dots$  be a G.P. with common ratio  $r$ .

$$\Rightarrow \frac{a_{n+1}}{a_n} = r \text{ for all } n \in \mathbb{N}$$

If each term of a G.P. is raised to the power  $x$ , we get the sequence

$$a_1^x, a_2^x, a_3^x, \dots, a_n^x, \dots$$

$$\text{Now, } \frac{(a_{n+1})^x}{(a_n)^x} = \left( \frac{a_{n+1}}{a_n} \right)^x = r^x \text{ for all } n \in \mathbb{N}$$

Hence,  $a_1^x, a_2^x, a_3^x, \dots, a_n^x, \dots$  is also a G.P.

**Question 7.**

If  $a, b$  and  $c$  are in A.P.  $a, x, b$  are in G.P. whereas  $b, y$  and  $c$  are also in G.P. Show that :  $x^2, b^2, y^2$  are in A.P.

**Solution:**

$a, b$  and  $c$  are in A.P.

$$\Rightarrow 2b = a + c$$

$a, x$  and  $b$  are in G.P.

$$\Rightarrow x^2 = ab$$

$b, y$  and  $c$  are in G.P.

$$\Rightarrow y^2 = bc$$

Now,

$$x^2 + y^2 = ab + bc$$

$$= b(a + c)$$

$$= b \times 2b$$

$$= 2b^2$$

$$\Rightarrow x^2, b^2 \text{ and } y^2 \text{ are in A.P.}$$

**Question 8.**

If  $a, b, c$  are in G.P. and  $a, x, b, y, c$  are in A.P., prove that :

$$(i) \frac{1}{x} + \frac{1}{y} = \frac{2}{b} \quad (ii) \frac{a}{x} + \frac{c}{y} = 2$$

**Solution 8(i).**

a, b and c are in G.P.

$$\Rightarrow b^2 = ac$$

a, x, b, y and c are in A.P.

$$\Rightarrow 2x = a + b \Rightarrow x = \frac{a+b}{2}$$

$$2b = x + y \Rightarrow b = \frac{x+y}{2}$$

$$2y = b + c \Rightarrow y = \frac{b+c}{2}$$

Now,

$$\begin{aligned} \frac{1}{x} + \frac{1}{y} &= \frac{2}{a+b} + \frac{2}{b+c} \\ &= \frac{2b + 2c + 2a + 2b}{ab + ac + b^2 + bc} \\ &= \frac{2a + 2c + 4b}{ab + b^2 + b^2 + bc} \\ &= \frac{2a + 2c + 4b}{ab + 2b^2 + bc} \\ &= \frac{2(a + c + 2b)}{b(a + 2b + c)} \\ &= \frac{2}{b} \end{aligned}$$

**Solution 8(ii).**

a, b and c are in G.P.

$$\Rightarrow b^2 = ac$$

a, x, b, y and c are in A.P.

$$\Rightarrow 2x = a + b \Rightarrow x = \frac{a+b}{2}$$

$$2b = x + y \Rightarrow b = \frac{x+y}{2}$$

$$2y = b + c \Rightarrow y = \frac{b+c}{2}$$

Now,

$$\begin{aligned}\frac{a}{x} + \frac{c}{y} &= \frac{2a}{a+b} + \frac{2c}{b+c} \\ &= \frac{2a(b+c) + 2c(a+b)}{(a+b)(b+c)} \\ &= \frac{2ab + 2ac + 2ac + 2bc}{ab + ac + b^2 + bc} \\ &= \frac{2ab + 4ac + 2bc}{ab + b^2 + b^2 + bc} \\ &= \frac{2(ab + 2ac + bc)}{ab + 2b^2 + bc} \\ &= \frac{2(ab + 2ac + bc)}{ab + 2ac + bc} \\ &= 2\end{aligned}$$

### Question 9.

If  $a$ ,  $b$  and  $c$  are in A.P. and also in G.P., show that:  $a = b = c$ .

#### Solution:

$a$ ,  $b$  and  $c$  are in A.P.

$$\Rightarrow 2b = a + c$$

$$\Rightarrow b = \frac{a+c}{2}$$

$a$ ,  $b$  and  $c$  are also in G.P.

$$\Rightarrow b^2 = ac$$

$$\Rightarrow \left(\frac{a+c}{2}\right)^2 = ac$$

$$\Rightarrow \frac{a^2 + c^2 + 2ac}{4} = ac$$

$$\Rightarrow a^2 + c^2 + 2ac = 4ac$$

$$\Rightarrow a^2 + c^2 - 2ac = 0$$

$$\Rightarrow (a-c)^2 = 0$$

$$\Rightarrow a - c = 0$$

$$\Rightarrow a = c$$

Now,  $2b = a + c$

$$\Rightarrow 2b = a + a$$

$$\Rightarrow 2b = 2a$$

$$\Rightarrow b = a$$

Thus, we have  $a = b = c$

**Question 10.**

The first term of a G.P. is  $a$  and its  $n^{\text{th}}$  term is  $b$ , where  $n$  is an even number. If the product of first  $n$  numbers of this G.P. is  $P$ ; prove that :  $p^2 = (ab)^n$ .

**Solution:**

For a G.P.,

First term =  $a$

Let the common ratio =  $r$

$n^{\text{th}}$  term =  $b$

$$\Rightarrow ar^{n-1} = b$$

$P$  = Product of first  $n$  numbers of the given G.P.

$$\Rightarrow P = a \times ar \times ar^2 \times ar^3 \times \dots \times ar^{n-1}$$

$$\Rightarrow P = a \times ar \times ar^2 \times ar^3 \times \dots \times b$$

$$\Rightarrow P = a \times ar \times ar^2 \times ar^3 \times \dots \times \frac{b}{r^2} \times \frac{b}{r} \times b$$

$$\Rightarrow P = (ab) \times \left( ar \times \frac{b}{r} \right) \times \left( ar^2 \times \frac{b}{r^2} \right) \times \dots \times \frac{n}{2} \text{ terms}$$

$$\Rightarrow P = (ab) \times (ab) \times (ab) \times \dots \times \frac{n}{2} \text{ terms}$$

$$\Rightarrow P = (ab)^{\frac{n}{2}}$$

$$\Rightarrow P = \sqrt{ab^n}$$

$$\Rightarrow p^2 = ab^n$$

**Question 11.**

If  $a, b, c$  and  $d$  are consecutive terms of a G.P. ; prove that :  $(a^2 + b^2), (b^2 + c^2)$  and  $(c^2 + d^2)$  are in GP.

**Solution:**

Let  $r$  be the common ratio of this G.P.

Given :  $a, b, c, d$  are in G.P.

$$\Rightarrow 1^{\text{st}} = a,$$

$$2^{\text{nd}} \text{ term} = b = ar,$$

$$3^{\text{rd}} \text{ term} = c = ar^2$$

$$4^{\text{th}} \text{ term} = d = ar^3$$

$$\begin{aligned}\text{Now, } (b^2 + c^2)^2 &= [(ar)^2 + (ar^2)^2]^2 \\ &= [a^2r^2 + a^2r^4]^2 \\ &= [a^2r^2(1 + r^2)]^2 \\ &= a^4r^4(1 + r^2)^2\end{aligned}$$

$$\begin{aligned}\text{And, } (a^2 + b^2) \times (c^2 + d^2) &= [a^2 + (ar)^2] \times [(ar^2)^2 + (ar^3)^2] \\ &= [a^2 + a^2r^2] \times [a^2r^4 + a^2r^6] \\ &= a^2(1 + r^2) \times a^2r^4(1 + r^2) \\ &= a^4r^4(1 + r^2)^2\end{aligned}$$

$$\Rightarrow (b^2 + c^2)^2 = (a^2 + b^2) \times (c^2 + d^2)$$

$$\text{i.e. } \frac{b^2 + c^2}{a^2 + b^2} = \frac{c^2 + d^2}{b^2 + c^2}$$

Hence,  $(a^2 + b^2)$ ,  $(b^2 + c^2)$  and  $(c^2 + d^2)$  are in G.P.

**Question 12.**

If  $a, b, c$  and  $d$  are consecutive terms of a G.P. To prove:

$\frac{1}{a^2 + b^2}$ ,  $\frac{1}{b^2 + c^2}$  and  $\frac{1}{c^2 + d^2}$  are in G.P.

**Solution:**

Let  $r$  be the common ratio of this G.P.

Given:  $a, b, c, d$  are in G.P.

$$\Rightarrow 1^{\text{st}} = a,$$

$$2^{\text{nd}} \text{ term} = b = ar,$$

$$3^{\text{rd}} \text{ term} = c = ar^2$$

$$4^{\text{th}} \text{ term} = d = ar^3$$

$$\begin{aligned} \text{Now, } \left( \frac{1}{b^2 + c^2} \right)^2 &= \left[ \frac{1}{(ar)^2 + (ar^2)^2} \right]^2 \\ &= \left[ \frac{1}{a^2r^2 + a^2r^4} \right]^2 \\ &= \frac{1}{a^4r^4} \left[ \frac{1}{1+r^2} \right]^2 \\ &= \frac{1}{a^4r^4} \times \frac{1}{(1+r^2)^2} \end{aligned}$$

$$\begin{aligned} \text{And, } \left( \frac{1}{a^2 + b^2} \right) \times \left( \frac{1}{c^2 + d^2} \right) &= \left[ \frac{1}{a^2 + (ar)^2} \right] \times \left[ \frac{1}{(ar^2)^2 + (ar^3)^2} \right] \\ &= \left[ \frac{1}{a^2 + a^2r^2} \right] \times \left[ \frac{1}{a^2r^4 + a^2r^6} \right] \\ &= \frac{1}{a^2} \left( \frac{1}{1+r^2} \right) \times \frac{1}{a^2r^4} \left( \frac{1}{1+r^2} \right) \\ &= \frac{1}{a^4r^4} \times \frac{1}{(1+r^2)^2} \end{aligned}$$

$$\Rightarrow \left( \frac{1}{b^2 + c^2} \right)^2 = \left( \frac{1}{a^2 + b^2} \right) \times \left( \frac{1}{c^2 + d^2} \right)$$

Hence,  $\frac{1}{a^2 + b^2}, \frac{1}{b^2 + c^2}$  and  $\frac{1}{c^2 + d^2}$  are in G.P.

**Exercise 11D****Question 1.**

Find the sum of G.P. :

(i)  $1 + 3 + 9 + 27 + \dots$  to 12 terms.

(ii)  $0.3 + 0.03 + 0.003 + 0.0003 + \dots$  to 8 terms.

(iii)  $1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots$  to 9 terms.

(iv)  $1 - \frac{1}{3} + \frac{1}{3^2} - \frac{1}{3^3} + \dots$  to  $n$  terms.

(v)  $\frac{x+y}{x-y} + 1 + \frac{x-y}{x+y} + \dots$  upto  $n$  terms.

(vi)  $\sqrt{3} + \frac{1}{\sqrt{3}} + \frac{1}{3\sqrt{3}} + \dots$  to  $n$  terms.

**Solution 1(i).**

Given GP. :  $1 + 3 + 9 + 27 + \dots$

Here,

first term,  $a = 1$

common ratio,  $r = \frac{3}{1} = 3$  ( $r > 1$ )

number of terms to be added,  $n = 12$

$$\therefore S_n = \frac{a(r^n - 1)}{r - 1}$$

$$\Rightarrow S_{12} = \frac{1(3^{12} - 1)}{3 - 1} = \frac{3^{12} - 1}{2} = \frac{531441 - 1}{2} = \frac{531440}{2} = 265720$$

**Solution 1(ii).**

Given GP. :  $0.3 + 0.03 + 0.003 + 0.0003 + \dots$

Here,

first term,  $a = 0.3$

common ratio,  $r = \frac{0.03}{0.3} = 0.1$  ( $r < 1$ )

number of terms to be added,  $n = 8$

$$\therefore S_n = \frac{a(1 - r^n)}{1 - r}$$

$$\Rightarrow S_8 = \frac{0.3(1 - (0.1)^8)}{1 - 0.1} = \frac{0.3(1 - (0.1)^8)}{0.9} = \frac{1 - (0.1)^8}{3} = \frac{1}{3} \left( 1 - \frac{1}{10^8} \right)$$

**Solution 1(iii).**

Given GP. :  $1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots$

Here,

first term,  $a = 1$

common ratio,  $r = \frac{-1/2}{1} = -\frac{1}{2}$  ( $r < 1$ )

number of terms to be added,  $n = 9$

$$\therefore S_n = \frac{a(1-r^n)}{1-r}$$

$$\Rightarrow S_8 = \frac{1 \left( 1 - \left( -\frac{1}{2} \right)^9 \right)}{1 - \left( -\frac{1}{2} \right)}$$

$$= \frac{1 - \left( -\frac{1}{2} \right)^9}{1 + \frac{1}{2}}$$

$$= \frac{1 + \frac{1}{2^9}}{\frac{3}{2}}$$

$$= \frac{2}{3} \left( 1 + \frac{1}{2^9} \right)$$

$$= \frac{2}{3} \left( 1 + \frac{1}{512} \right)$$

$$= \frac{2}{3} \times \frac{513}{512}$$

$$= \frac{171}{256}$$

**Solution 1(iv).**

Given G.P.:  $1 - \frac{1}{3} + \frac{1}{3^2} - \frac{1}{3^3} + \dots$  upto n terms

Here,

first term,  $a = 1$

common ratio,  $r = \frac{-1/3}{1} = -\frac{1}{3}$  ( $r < 1$ )

number of terms to be added = n

$$\therefore S_n = \frac{a(1-r^n)}{1-r}$$

$$\Rightarrow S_n = \frac{1 \left( 1 - \left( -\frac{1}{3} \right)^n \right)}{1 - \left( -\frac{1}{3} \right)}$$

$$= \frac{1 \left( 1 - \left( -\frac{1}{3} \right)^n \right)}{1 + \frac{1}{3}}$$

$$= \frac{\left[ 1 - \left( -\frac{1}{3} \right)^n \right]}{\frac{4}{3}}$$

$$= \frac{3}{4} \left[ 1 - \left( -\frac{1}{3} \right)^n \right]$$

**Solution 1(v).**

Given G.P.:  $\frac{x+y}{x-y} + 1 + \frac{x-y}{x+y} + \dots$  upto n terms

Here,

$$\text{first term, } a = \frac{x+y}{x-y}$$

$$\text{common ratio, } r = \frac{1}{\frac{x+y}{x-y}} = \frac{x-y}{x+y} \quad (r < 1)$$

number of terms to be added = n

$$\therefore S_n = \frac{a(1-r^n)}{1-r}$$

$$\Rightarrow S_n = \frac{\frac{x+y}{x-y} \left( 1 - \left( \frac{x-y}{x+y} \right)^n \right)}{1 - \left( \frac{x-y}{x+y} \right)}$$

$$= \frac{\frac{x+y}{x-y} \left( 1 - \left( \frac{x-y}{x+y} \right)^n \right)}{\frac{x+y - x+y}{x+y}}$$

$$= \frac{\frac{x+y}{x-y} \left( 1 - \left( \frac{x-y}{x+y} \right)^n \right)}{\frac{2y}{x+y}}$$

$$= \frac{(x+y)^2 \left( 1 - \left( \frac{x-y}{x+y} \right)^n \right)}{2y(x-y)}$$

**Solution 1(vi).**

Given G.P.:  $\sqrt{3} + \frac{1}{\sqrt{3}} + \frac{1}{3\sqrt{3}} + \dots$  upto n terms

Here,

first term,  $a = \sqrt{3}$

common ratio,  $r = \frac{\frac{1}{\sqrt{3}}}{\sqrt{3}} = \frac{1}{3}$  ( $r < 1$ )

number of terms to be added = n

$$\therefore S_n = \frac{a(1-r^n)}{1-r}$$

$$\Rightarrow S_n = \frac{\sqrt{3} \left( 1 - \left( \frac{1}{3} \right)^n \right)}{1 - \frac{1}{3}}$$

$$= \frac{\sqrt{3} \left( 1 - \frac{1}{3^n} \right)}{\frac{2}{3}}$$

$$= \frac{3\sqrt{3}}{2} \left( 1 - \frac{1}{3^n} \right)$$

**Question 2.**

How many terms of the geometric progression  $1 + 4 + 16 + 64 + \dots$  must be added to get sum equal to 5461?

**Solution:**

Given G.P.:  $1 + 4 + 16 + 64 + \dots$

Here,

first term,  $a = 1$

common ratio,  $r = \frac{4}{1} = 4$  ( $r > 1$ )

Let the number of terms to be added = n

Then,  $S_n = 5461$

$$\Rightarrow \frac{a(r^n - 1)}{r - 1} = 5461$$

$$\Rightarrow \frac{1(4^n - 1)}{4 - 1} = 5461$$

$$\Rightarrow \frac{4^n - 1}{3} = 5461$$

$$\Rightarrow 4^n - 1 = 16383$$

$$\Rightarrow 4^n = 16384$$

$$\Rightarrow 4^n = 4^7$$

$$\Rightarrow n = 7$$

Hence, required number of terms = 7

### Question 3.

The first term of a G.P. is 27 and its 8<sup>th</sup> term is  $\frac{1}{81}$ . Find the sum of its first 10 terms.

#### Solution:

Given,

First term,  $a = 27$

$$8^{\text{th}} \text{ term} = ar^7 = \frac{1}{81}$$

$n = 10$

Now,

$$\frac{ar^7}{a} = \frac{1/81}{27}$$

$$\Rightarrow r^7 = \frac{1}{2187}$$

$$\Rightarrow r^7 = \left(\frac{1}{3}\right)^7$$

$$\Rightarrow r = \frac{1}{3} \quad (r < 1)$$

$$\therefore S_n = \frac{a(1 - r^n)}{1 - r}$$

$$\Rightarrow S_{10} = \frac{27 \left(1 - \left(\frac{1}{3}\right)^{10}\right)}{1 - \frac{1}{3}}$$

$$= \frac{27 \left(1 - \frac{1}{3^{10}}\right)}{\frac{2}{3}}$$

$$\frac{81}{2} \left(1 - \frac{1}{3^{10}}\right)$$

**Question 4.**

A boy spends ₹ 10 on first day, ₹ 20 on second day, ₹ 40 on third day and so on. Find how much, in all, will he spend in 12 days?

**Solution:**

Amount spent on 1<sup>st</sup> day = Rs. 10

Amount spent on 2<sup>nd</sup> day = Rs. 20

Amount spent on 3<sup>d</sup> day = Rs. 40 and so on

$$\text{Now, } \frac{20}{10} = 2, \frac{40}{20} = 2,$$

Thus, 10, 20, 40, ..... is a G.P. with first term,  $a = 10$   
and common ratio,  $r = 2$  ( $r > 1$ )

∴ Total amount spent in 12 days =  $S_{12}$

$$S_n = \frac{a(r^n - 1)}{r - 1}$$

$$\Rightarrow S_{12} = \frac{10(2^{12} - 1)}{2 - 1} = 10(2^{12} - 1) = 10(4096 - 1) = 10 \times 4095 = 40950$$

Hence, the total amount spent in 12 days is Rs. 40950.

**Question 5.**

The 4th and the 7th terms of a G.P. are  $\frac{1}{27}$  and  $\frac{1}{729}$  respectively. Find the sum of n terms of this G.P.

**Solution:**

For a G.P.,

$$4^{\text{th}} \text{ term} = ar^3 = \frac{1}{27}$$

$$7^{\text{th}} \text{ term} = ar^6 = \frac{1}{729}$$

$$\text{Now, } \frac{ar^6}{ar^3} = \frac{\frac{1}{729}}{\frac{1}{27}}$$

$$\Rightarrow r^3 = \frac{1}{27} = \left(\frac{1}{3}\right)^3$$

$$\Rightarrow r = \frac{1}{3} \quad (r < 1)$$

$$\Rightarrow a \times \frac{1}{27} = \frac{1}{27}$$

$$\Rightarrow a = 1$$

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$\Rightarrow S_n = \frac{1 \left( 1 - \left( \frac{1}{3} \right)^n \right)}{1 - \frac{1}{3}}$$

$$= \frac{\left( 1 - \frac{1}{3^n} \right)}{\frac{2}{3}}$$

$$= \frac{3}{2} \left( 1 - \frac{1}{3^n} \right)$$

### Question 6.

A geometric progression has common ratio = 3 and last term = 486. If the sum of its terms is 728 ; find its first term.

### Solution:

For a G.P.,

Common ratio,  $r = 3$  ( $r > 1$ )

Last term,  $l = 486$

$S = 728$

$$\Rightarrow \frac{l r - a}{r - 1} = 728$$

$$\Rightarrow \frac{486 \times 3 - a}{3 - 1} = 728$$

$$\Rightarrow \frac{1458 - a}{2} = 728$$

$$\Rightarrow 1458 - a = 1456$$

Hence, the first term is 2.

**Question 7.**

Find the sum of G.P. : 3, 6, 12, ..... 1536.

**Solution:**

Given G.P. : 3, 6, 12, ....., 1536

Here,

First term,  $a = 3$

Common ratio,  $r = \frac{6}{3} = 2$  ( $r > 1$ )

Last term,  $l = 1536$

$$\begin{aligned}\therefore \text{ Required sum} &= \frac{l r - a}{r - 1} \\ &= \frac{1536 \times 2 - 3}{2 - 1} \\ &= 3072 - 3 \\ &= 3069\end{aligned}$$

**Question 8.**

How many terms of the series  $2 + 6 + 18 + \dots$  must be taken to make the sum equal to 728 ?

**Solution:**

Given series:  $2 + 6 + 18 + \dots$

$$\text{Now, } \frac{6}{2} = 3, \frac{18}{6} = 3$$

Thus, given series is a G.P. with first term,  $a = 2$   
and common ratio,  $r = 3$  ( $r > 1$ )

Let the number of terms to be added =  $n$

Then,  $S_n = 728$

$$\Rightarrow \frac{a(r^n - 1)}{r - 1} = 728$$

$$\Rightarrow \frac{2(3^n - 1)}{3 - 1} = 728$$

$$\Rightarrow 3^n - 1 = 728$$

$$\Rightarrow 3^n = 729$$

$$\Rightarrow 3^n = 3^6$$

$$\Rightarrow n = 6$$

Hence, required number of terms = 6

**Question 9.**

In a G.P., the ratio between the sum of first three terms and that of the first six terms is 125 : 152.

Find its common ratio.

**Solution:**

Let  $a$  be the first term and  $r$  be the common ratio of given G.P.

$$\text{Now, sum of first three terms} = S_3 = \frac{a(r^3 - 1)}{r - 1}$$

$$\text{Now, sum of first six terms} = S_6 = \frac{a(r^6 - 1)}{r - 1}$$

It is given that

$$\frac{\frac{a(r^3 - 1)}{r - 1}}{\frac{a(r^6 - 1)}{r - 1}} = \frac{125}{152}$$

$$\Rightarrow \frac{r^3 - 1}{r^6 - 1} = \frac{125}{152}$$

$$\Rightarrow \frac{r^3 - 1}{(r^3)^2 - (1)^2} = \frac{125}{152}$$

$$\Rightarrow \frac{r^3 - 1}{(r^3 - 1)(r^3 + 1)} = \frac{125}{152}$$

$$\Rightarrow \frac{1}{r^3 + 1} = \frac{125}{152}$$

$$\Rightarrow r^3 + 1 = \frac{152}{125}$$

$$\Rightarrow r^3 = \frac{152}{125} - 1 = \frac{152 - 125}{125} = \frac{27}{125}$$

$$\Rightarrow r^3 = \left(\frac{3}{5}\right)^3$$

$$\Rightarrow r = \frac{3}{5}$$

Hence, the common ratio is  $\frac{3}{5}$ .

**Question 10.**

Find how many terms of G.P.  $\frac{2}{9} - \frac{1}{3} + \frac{1}{2} \dots\dots\dots$  must be added to get the sum equal to  $\frac{55}{72}$ ?

**Solution:**

Given G.P.:  $\frac{2}{9} - \frac{1}{3} + \frac{1}{2} \dots\dots\dots$

Here,

First term,  $a = \frac{2}{9}$

Common ratio,  $r = \frac{-1/3}{2/9} = -\frac{3}{2} < 1$

Let required number of terms be  $n$ .

$$\Rightarrow S_n = \frac{55}{72}$$

$$\Rightarrow \frac{a(1-r^n)}{1-r} = \frac{55}{72}$$

$$\Rightarrow \frac{\frac{2}{9} \left( 1 - \left( \frac{-3}{2} \right)^n \right)}{1 - \left( \frac{-3}{2} \right)} = \frac{55}{72}$$

$$\Rightarrow \frac{\frac{2}{9} \left( 1 - \left( \frac{-3}{2} \right)^n \right)}{\frac{5}{2}} = \frac{55}{72}$$

$$\Rightarrow \frac{2}{9} \left( 1 - \left( \frac{-3}{2} \right)^n \right) = \frac{55}{72} \times \frac{5}{2}$$

$$\Rightarrow 1 - \left( \frac{-3}{2} \right)^n = \frac{55}{72} \times \frac{5}{2} \times \frac{9}{2}$$

$$\Rightarrow 1 - \left( \frac{-3}{2} \right)^n = \frac{275}{32}$$

$$\Rightarrow 1 - \frac{275}{32} = \left( \frac{-3}{2} \right)^n$$

$$\Rightarrow -\frac{243}{32} = \left( \frac{-3}{2} \right)^n$$

$$\Rightarrow \left( \frac{-3}{2} \right)^5 = \left( \frac{-3}{2} \right)^n$$

$$\Rightarrow n = 5$$

$\therefore$  Required number of terms = 5

**Question 11.**

If the sum  $1 + 2 + 2^2 + \dots + 2^{n-1}$  is 255, find the value of  $n$ .

**Solution:**

Required series:  $1 + 2 + 2^2 + \dots + 2^{n-1}$

$$\text{Now, } \frac{2}{1} = 2, \frac{2^2}{2} = 2$$

Thus, given series is a G.P. with first term,  $a = 1$

common ratio,  $r = 2$  ( $r > 1$ )

Last term,  $l = 2^{n-1}$

Let there be  $n$  terms in the series.

Then,  $S_n = 255$

$$\Rightarrow \frac{lr - a}{r - 1} = 255$$

$$\Rightarrow \frac{2^{n-1} \times 2 - 1}{2 - 1} = 255$$

$$\Rightarrow 2^{n-1} \times 2 - 1 = 255$$

$$\Rightarrow 2^{n-1} \times 2 = 256$$

$$\Rightarrow 2^{n-1} = 128$$

$$\Rightarrow 2^{n-1} = 2^7$$

$$\Rightarrow n - 1 = 7$$

$$\Rightarrow n = 8$$

**Question 12.**

Find the geometric mean between :

(i)  $\frac{4}{9}$  and  $\frac{9}{4}$

(ii) 14 and  $\frac{7}{32}$

(iii)  $2a$  and  $8a^3$

**Solution 12(i).**

$$\text{Geometric mean between } \frac{4}{9} \text{ and } \frac{9}{4} = \sqrt{\frac{4}{9} \times \frac{9}{4}} = \sqrt{1} = 1$$

**Solution 12(ii).**

$$\text{Geometric mean between } 14 \text{ and } \frac{7}{32} = \sqrt{14 \times \frac{7}{32}} = \sqrt{\frac{49}{16}} = \frac{7}{4} = 1\frac{3}{4}$$

**Solution 12(iii).**

Geometric mean between  $2a$  and  $8a^3 = \sqrt{2a \times 8a^3} = \sqrt{16 \times a^4} = 4a^2$

**Question 13.**

The sum of three numbers in G.P. is  $\frac{39}{10}$  and their product is 1. Find the numbers.

**Solution:**

Let the numbers be  $\frac{a}{r}$ ,  $a$  and  $ar$ .

$$\Rightarrow \frac{a}{r} \times a \times ar = 1$$

$$\Rightarrow a^3 = 1$$

$$\Rightarrow a = 1$$

$$\text{Now, } \frac{a}{r} + a + ar = \frac{39}{10}$$

$$\Rightarrow \frac{1}{r} + 1 + r = \frac{39}{10}$$

$$\Rightarrow \frac{1+r+r^2}{r} = \frac{39}{10}$$

$$\Rightarrow 10 + 10r + 10r^2 = 39r$$

$$\Rightarrow 10r^2 - 29r + 10 = 0$$

$$\Rightarrow 10r^2 - 25r - 4r + 10 = 0$$

$$\Rightarrow 5r(2r - 5) - 2(2r - 5) = 0$$

$$\Rightarrow (2r - 5)(5r - 2) = 0$$

$$\Rightarrow r = \frac{5}{2} \text{ or } r = \frac{2}{5}$$

Thus, required terms are:

$$\begin{aligned} \frac{a}{r}, a, ar &= \frac{1}{\frac{5}{2}}, 1, 1 \times \frac{5}{2} \quad \text{OR} \quad \frac{1}{\frac{2}{5}}, 1, 1 \times \frac{2}{5} \\ &= \frac{2}{5}, 1, \frac{5}{2} \quad \text{OR} \quad \frac{5}{2}, 1, \frac{2}{5} \end{aligned}$$

**Question 14.**

The first term of a G.P. is  $-3$  and the square of the second term is equal to its 4<sup>th</sup> term. Find its 7<sup>th</sup> term.

**Solution:**

For a G.P.,

First term,  $a = -3$

It is given that,

$$(2^{\text{nd}} \text{ term})^2 = 4^{\text{th}} \text{ term}$$

$$\Rightarrow (ar)^2 = ar^3$$

$$\Rightarrow a^2r^2 = ar^3$$

$$\Rightarrow a = r$$

$$\Rightarrow r = -3$$

$$\text{Now, } 7^{\text{th}} \text{ term} = ar^6 = -3 \times (-3)^6 = -3 \times 729 = -2187$$

### Question 15.

Find the 5<sup>th</sup> term of the G.P.  $\frac{5}{2}, 1, \dots$

**Solution:**

$$\text{First term (a)} = \frac{5}{2}$$

$$\text{And, common ratio (r)} = \frac{1}{\frac{5}{2}} = \frac{2}{5}$$

$$\text{Now, } t_n = ar^{n-1}$$

$$\Rightarrow 5^{\text{th}} \text{ term} = t_5 = \frac{5}{2} \times \left(\frac{5}{2}\right)^{5-1} = \frac{5}{2} \times \left(\frac{2}{5}\right)^4 = \left(\frac{2}{5}\right)^3 = \frac{8}{125}$$

### Question 16.

The first two terms of a G.P. are 125 and 25 respectively. Find the 5th and the 6th terms of the G.P.

**Solution:**

$$\text{First term (a)} = 125$$

$$\text{And, common ratio (r)} = \frac{25}{125} = \frac{1}{5}$$

$$\text{Now, } t_n = ar^{n-1}$$

$$\Rightarrow 5^{\text{th}} \text{ term} = t_5 = 125 \times \left(\frac{1}{5}\right)^{5-1} = 125 \times \left(\frac{1}{5}\right)^4 = 125 \times \frac{1}{625} = \frac{1}{5}$$

$$\Rightarrow 6^{\text{th}} \text{ term} = t_6 = 125 \times \left(\frac{1}{5}\right)^{6-1} = 125 \times \left(\frac{1}{5}\right)^5 = 125 \times \frac{1}{3125} = \frac{1}{25}$$

**Question 17.**

Find the sum of the sequence  $-\frac{1}{3}, 1, -3, 9, \dots$  upto 8 terms.

**Solution:**

Here,

$$\frac{1}{-\frac{1}{3}} = \frac{-3}{1} = \frac{9}{-3} = -3$$

Thus, the given sequence is a G.P. with first term  $(a) = -\frac{1}{3}$  and common ratio  $(r) = -3$  ( $r < 1$ ).

Number of terms to be added,  $n = 8$

$$\therefore S_n = \frac{a(1-r^n)}{1-r}$$

$$\Rightarrow S_8 = \frac{-\frac{1}{3}(1-(-3)^8)}{1+3} = \frac{-1+3^8}{12} = \frac{1}{12}(3^8 - 1)$$

**Question 18.**

The first term of a G.P. is 27. If the 8th term be  $\frac{1}{81}$ , what will be the sum of 10 terms?

**Solution:**

Given,

First term,  $a = 27$

$$8^{\text{th}} \text{ term} = ar^7 = \frac{1}{81}$$

$n = 10$

Now,

$$\frac{ar^7}{a} = \frac{1/81}{27}$$

$$\Rightarrow r^7 = \frac{1}{2187}$$

$$\Rightarrow r^7 = \left(\frac{1}{3}\right)^7$$

$$\Rightarrow r = \frac{1}{3} \quad (r < 1)$$

$$\therefore S_n = \frac{a(1-r^n)}{1-r}$$

$$\begin{aligned}
\Rightarrow S_{10} &= \frac{27\left(1 - \left(\frac{1}{3}\right)^{10}\right)}{1 - \frac{1}{3}} \\
&= \frac{27\left(1 - \frac{1}{3^{10}}\right)}{\frac{2}{3}} \\
&= \frac{81}{2}\left(1 - \frac{1}{3^{10}}\right) \\
&= \frac{81}{2}(1 - 3^{-10})
\end{aligned}$$

**Question 19.**

Find a G.P. for which the sum of first two terms is -4 and the fifth term is 4 times the third term.

**Solution:**

Let the five terms of the given G.P. be

$$\frac{a}{r^2}, \frac{a}{r}, a, ar, ar^2$$

Given, sum of first two terms = -4

$$\frac{a}{r^2} + \frac{a}{r} = -4$$

$$\Rightarrow \frac{a + ar}{r^2} = -4$$

$$\Rightarrow a + ar = -4r^2$$

$$\Rightarrow a(1 + r) = -4r^2$$

$$\Rightarrow a = -\frac{4r^2}{1+r}$$

And, 5<sup>th</sup> term = 4(3<sup>rd</sup> term)

$$\Rightarrow ar^2 = 4(a)$$

$$\Rightarrow r^2 = 4$$

$$\Rightarrow r = \pm 2$$

When  $r = +2$ ,

$$a = -\frac{4(2)^2}{1+2} = -\frac{16}{3}$$

When  $r = -2$ ,

$$a = -\frac{4(-2)^2}{1-2} = 16$$

Thus, the required terms are  $\frac{a}{r^2}, \frac{a}{r}, a, ar, ar^2$ .

$$\text{i.e. } \frac{-16}{4}, \frac{-16}{2}, -\frac{16}{3}, -\frac{16}{3} \times 2, -\frac{16}{3} \times 4 \text{ OR } \frac{16}{4}, \frac{16}{-2}, 16, 16(-2), 16 \times 4$$

$$\text{i.e. } -\frac{4}{3}, -\frac{8}{3}, -\frac{16}{3}, -\frac{32}{3}, -\frac{64}{3} \text{ OR } 4, -8, 16, -32, 64$$

## Additional Questions

### Question 1.

Find the sum of  $n$  terms of the series :

(i)  $4 + 44 + 444 + \dots$

(ii)  $0.8 + 0.88 + 0.888 + \dots$

### Solution 1(i).

$$\begin{aligned} \text{Required sum} &= 4 + 44 + 444 + \dots \text{ upto } n \text{ terms} \\ &= 4(1 + 11 + 111 + \dots \text{ upto } n \text{ terms}) \\ &= \frac{4}{9}(9 + 99 + 999 + \dots \text{ upto } n \text{ terms}) \\ &= \frac{4}{9}[(10 - 1) + (100 - 1) + (1000 - 1) + \dots \text{ upto } n \text{ terms}] \\ &= \frac{4}{9} \left[ (10 + 10^2 + 10^3 + \dots \text{ upto } n \text{ terms}) \right. \\ &\quad \left. - (1 + 1 + 1 + \dots \text{ upto } n \text{ terms}) \right] \\ &= \frac{4}{9} \left[ \frac{10(10^n - 1)}{10 - 1} - n \right] \\ &= \frac{4}{9} \left[ \frac{10}{9}(10^n - 1) - n \right] \end{aligned}$$

**Solution 1(ii).**

$$\begin{aligned}
\text{Required sum} &= 0.8 + 0.88 + 0.888 + \dots \text{ upto } n \text{ terms} \\
&= 8(0.1 + 0.11 + 0.111 + \dots \text{ upto } n \text{ terms}) \\
&= \frac{8}{9}(0.9 + 0.99 + 0.999 + \dots \text{ upto } n \text{ terms}) \\
&= \frac{8}{9}[(1 - 0.1) + (1 - 0.01) + (1 - 0.001) + \dots \text{ upto } n \text{ terms}] \\
&= \frac{8}{9} \left[ \begin{array}{l} (1 + 1 + 1 + \dots \text{ upto } n \text{ terms}) \\ - (0.1 + 0.01 + 0.001 + \dots \text{ upto } n \text{ terms}) \end{array} \right] \\
&= \frac{8}{9} \left[ \begin{array}{l} (1 + 1 + 1 + \dots \text{ upto } n \text{ terms}) \\ - \left( \frac{1}{10} + \frac{1}{100} + \frac{1}{1000} + \dots \text{ upto } n \text{ terms} \right) \end{array} \right] \\
&= \frac{8}{9} \left[ n - \frac{\frac{1}{10} \left( 1 - \frac{1}{10^n} \right)}{1 - \frac{1}{10}} \right] \quad \left[ \because r = \frac{1}{10} < 1 \right] \\
&= \frac{8}{9} \left[ n - \frac{10}{9} \times \frac{1}{10} \left( 1 - \frac{1}{10^n} \right) \right] \\
&= \frac{8}{9} \left[ n - \frac{1}{9} \left( 1 - \frac{1}{10^n} \right) \right]
\end{aligned}$$

**Question 2.**

Find the sum of infinite terms of each of the following geometric progression:

(i)  $1 + \frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \dots$

(ii)  $1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots$

(iii)  $\frac{1}{3} + \frac{1}{3^2} - \frac{1}{3^3} + \dots$

(iv)  $\sqrt{2} - \frac{1}{\sqrt{2}} + \frac{1}{2\sqrt{2}} - \frac{1}{4\sqrt{2}} + \dots$

(v)  $\sqrt{3} + \frac{1}{\sqrt{3}} + \frac{1}{3\sqrt{3}} + \frac{1}{9\sqrt{3}} + \dots$

**Solution 2(i).**

$$\text{Given GP. : } 1 + \frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \dots$$

Here,

$$\text{First term, } a = 1$$

$$\text{Common ratio, } r = \frac{1/3}{1} = \frac{1}{3} \left( |r| = \left| \frac{1}{3} \right| = \frac{1}{3} < 1 \right)$$

$$\therefore \text{ Required sum} = \frac{a}{1-r} = \frac{1}{1-\frac{1}{3}} = \frac{3}{2}$$

**Solution 2(ii).**

$$\text{Given GP. : } 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots$$

Here,

$$\text{First term, } a = 1$$

$$\text{Common ratio, } r = \frac{-1/2}{1} = -\frac{1}{2} \left( |r| = \left| -\frac{1}{2} \right| = \frac{1}{2} < 1 \right)$$

$$\therefore \text{ Required sum} = \frac{a}{1-r} = \frac{1}{1-\left(-\frac{1}{2}\right)} = \frac{1}{1+\frac{1}{2}} = \frac{2}{3}$$

**Solution 2(iii).**

$$\text{Given GP. : } \frac{1}{3} + \frac{1}{3^2} + \frac{1}{3^3} + \dots$$

Here,

$$\text{First term, } a = \frac{1}{3}$$

$$\text{Common ratio, } r = \frac{1/3^2}{1/3} = \frac{1}{3} \left( |r| = \left| \frac{1}{3} \right| = \frac{1}{3} < 1 \right)$$

$$\therefore \text{ Required sum} = \frac{a}{1-r} = \frac{\frac{1}{3}}{1-\frac{1}{3}} = \frac{\frac{1}{3}}{\frac{2}{3}} = \frac{1}{2}$$

**Solution 2(iv).**

$$\text{Given GP. : } \sqrt{2} - \frac{1}{\sqrt{2}} + \frac{1}{2\sqrt{2}} - \frac{1}{4\sqrt{2}} + \dots$$

Here,

$$\text{First term, } a = \sqrt{2}$$

$$\text{Common ratio, } r = \frac{-1/\sqrt{2}}{\sqrt{2}} = -\frac{1}{2} \left( |r| = \left| -\frac{1}{2} \right| = \frac{1}{2} < 1 \right)$$

$$\therefore \text{ Required sum} = \frac{a}{1-r} = \frac{\sqrt{2}}{1 - \left(-\frac{1}{2}\right)} = \frac{\sqrt{2}}{1 + \frac{1}{2}} = \frac{2\sqrt{2}}{3}$$

**Solution 2(v).**

$$\text{Given GP. : } \sqrt{3} + \frac{1}{\sqrt{3}} + \frac{1}{3\sqrt{3}} + \frac{1}{9\sqrt{3}} + \dots$$

Here,

$$\text{First term, } a = \sqrt{3}$$

$$\text{Common ratio, } r = \frac{1/\sqrt{3}}{\sqrt{3}} = \frac{1}{3} \left( |r| = \left| \frac{1}{3} \right| = \frac{1}{3} < 1 \right)$$

$$\therefore \text{ Required sum} = \frac{a}{1-r} = \frac{\sqrt{3}}{1 - \frac{1}{3}} = \frac{\sqrt{3}}{\frac{2}{3}} = \frac{3\sqrt{3}}{2}$$

**Question 3.**

The second term of a G.P. is 9 and sum of its infinite terms is 48. Find its first three terms.

**Solution:**

Let  $a$  be the first term and  $r$  be the common ratio of a G.P.

$$2^{\text{nd}} \text{ term, } t_2 = ar = 9$$

$$\Rightarrow r = \frac{9}{a}$$

Sum of its infinite terms,  $S = 48$

$$\Rightarrow \frac{a}{1-r} = 48$$

$$\Rightarrow \frac{a}{1 - \frac{9}{a}} = 48$$

$$\Rightarrow \frac{a^2}{a-9} = 48$$

$$\Rightarrow a^2 = 48a - 432$$

$$\Rightarrow a^2 - 48a + 432 = 0$$

$$\Rightarrow a^2 - 36a - 12a + 432 = 0$$

$$\Rightarrow a(a-36) - 12(a-36) = 0$$

$$\Rightarrow (a-36)(a-12) = 0$$

$$\Rightarrow a = 36 \text{ or } a = 12$$

$$\text{When } a = 36, r = \frac{9}{36} = \frac{1}{4}$$

$$\Rightarrow 1^{\text{st}} \text{ term} = 36,$$

$$2^{\text{nd}} \text{ term} = ar = 36 \times \frac{1}{4} = 9$$

$$3^{\text{rd}} \text{ term} = ar^2 = 36 \times \frac{1}{16} = \frac{9}{4}$$

$$\text{When } a = 12, r = \frac{9}{12} = \frac{3}{4}$$

$$\Rightarrow 1^{\text{st}} \text{ term} = 12,$$

$$2^{\text{nd}} \text{ term} = ar = 12 \times \frac{3}{4} = 9$$

$$3^{\text{rd}} \text{ term} = ar^2 = 12 \times \frac{9}{16} = \frac{27}{4}$$

**Question 4.**

Find three geometric means between  $\frac{1}{3}$  and 432.

**Solution:**

Let  $G_1, G_2, G_3$  be three geometric means between  $a = \frac{1}{3}$  and  $b = 432$ .

Then,  $\frac{1}{3}, G_1, G_2, G_3, 432$  is a G.P.

Thus, we have

$$\text{First term} = a = \frac{1}{3}$$

$$5^{\text{th}} \text{ term of the G.P.} = ar^4 = 432$$

$$\Rightarrow \frac{1}{3} \times r^4 = 432$$

$$\Rightarrow r^4 = 1296$$

$$\Rightarrow r^4 = 6^4$$

$$\Rightarrow r = 6$$

$$\therefore G_1 = ar = \frac{1}{3} \times 6 = 2$$

$$G_2 = ar^2 = \frac{1}{3} \times 6 \times 6 = 12$$

$$G_3 = ar^3 = \frac{1}{3} \times 6 \times 6 \times 6 = 72$$

### Question 5.

Find :

- (i) two geometric means between 2 and 16
- (ii) four geometric means between 3 and 96.
- (iii) five geometric means between  $3\frac{5}{9}$  and  $40\frac{1}{2}$

### Solution 5(i).

Let  $G_1, G_2$  be two geometric means between  $a = 2$  and  $b = 16$ .

Then,  $2, G_1, G_2, 16$  is a G.P.

Thus, we have

$$\text{First term} = a = 2$$

$$4^{\text{th}} \text{ term of the G.P.} = ar^3 = 16$$

$$\Rightarrow 2 \times r^3 = 16$$

$$\Rightarrow r^3 = 8$$

$$\Rightarrow r^3 = 2^3$$

$$\Rightarrow r = 2$$

$$\therefore G_1 = ar = 2 \times 2 = 4$$

$$G_2 = ar^2 = 2 \times 2 \times 2 = 8$$

**Solution 5(ii).**

Let  $G_1, G_2, G_3, G_4$  be four geometric means between  $a = 3$  and  $b = 96$ .

Then,  $3, G_1, G_2, G_3, G_4, 96$  is a G.P.

Thus, we have

First term =  $a = 3$

6<sup>th</sup> term of the G.P. =  $ar^5 = 96$

$$\Rightarrow 3 \times r^5 = 96$$

$$\Rightarrow r^5 = 32$$

$$\Rightarrow r^5 = 2^5$$

$$\Rightarrow r = 2$$

$$\therefore G_1 = ar = 3 \times 2 = 6$$

$$G_2 = ar^2 = 3 \times 4 = 12$$

$$G_3 = ar^3 = 3 \times 8 = 24$$

$$G_4 = ar^4 = 3 \times 16 = 48$$

**Solution 5(iii).**

Let  $G_1, G_2, G_3, G_4, G_5$  be five geometric means between

$$a = 3\frac{5}{9} = \frac{32}{9} \text{ and } b = 40\frac{1}{2} = \frac{81}{2}.$$

Then,  $\frac{32}{9}, G_1, G_2, G_3, G_4, G_5, \frac{81}{2}$  is a G.P.

Thus, we have

$$\text{First term} = a = \frac{32}{9}$$

$$7^{\text{th}} \text{ term of the G.P.} = ar^6 = \frac{81}{2}$$

$$\Rightarrow \frac{32}{9} \times r^6 = \frac{81}{2}$$

$$\Rightarrow r^6 = \frac{81}{2} \times \frac{9}{32}$$

$$\Rightarrow r^6 = \frac{729}{64}$$

$$\Rightarrow r^6 = \left(\frac{3}{2}\right)^6$$

$$\Rightarrow r = \frac{3}{2}$$

$$\therefore G_1 = ar = \frac{32}{9} \times \frac{3}{2} = \frac{16}{3}$$

$$G_2 = ar^2 = \frac{32}{9} \times \frac{9}{4} = 8$$

$$G_3 = ar^3 = \frac{32}{9} \times \frac{27}{8} = 12$$

$$G_4 = ar^4 = \frac{32}{9} \times \frac{81}{16} = 18$$

$$G_5 = ar^5 = \frac{32}{9} \times \frac{243}{32} = 27$$

**Question 6.**

The sum of three numbers in G.P. is  $\frac{39}{10}$  and their product is 1. Find the numbers.

**Solution:**

Sum of three numbers in G.P. =  $\frac{39}{10}$  and their product = 1

Let number be  $\frac{a}{r}$ ,  $a$ ,  $ar$ , then

$$\frac{a}{r} \times a \times ar = 1 \Rightarrow a^3 = 1 = (1)^3$$

$$\therefore a = 1$$

$$\text{and } \frac{a}{r} + a + ar = \frac{39}{10}$$

$$\Rightarrow a \left( \frac{1}{r} + 1 + r \right) = \frac{39}{10}$$

$$\frac{1}{r} + 1 + r = \frac{39}{10} \times 1 = \frac{39}{10}$$

$$r + \frac{1}{r} = \frac{39}{10} - 1 = \frac{39-10}{10} = \frac{29}{10}$$

$$r^2 + 1 = \frac{29}{10}r$$

$$\begin{aligned}
10r^2 + 10 &= 29r \Rightarrow 10r^2 - 29r + 10 = 0 \\
\Rightarrow 10r^2 - 4r - 25r + 10 &= 0 \\
\Rightarrow 2r(5r - 2) - 5(5r - 2) &= 0 \\
\Rightarrow (5r - 2)(2r - 5) &= 0
\end{aligned}$$

$$\text{Either } 5r - 2 = 0, \text{ then } r = \frac{2}{5}$$

$$\text{or } 2r - 5 = 0, \text{ then } r = \frac{5}{2}$$

$$\therefore \text{ Numbers are } \frac{2}{5}, 1, \frac{4}{25}, \text{ or } \frac{5}{2}, 1, \frac{25}{4}$$

### Question 7.

Find the numbers in G.P. whose sum is 52 and the sum of whose product in pairs is 624.

### Solution:

Let the numbers be  $a$ ,  $ar$  and  $ar^2$ .

$$\Rightarrow a + ar + ar^2 = 52 \quad \dots(i)$$

$$\text{And, } (a \times ar) + (ar \times ar^2) + (ar^2 \times a) = 624$$

$$\Rightarrow a^2r + a^2r^3 + a^2r^2 = 624$$

$$\Rightarrow ar(a + ar^2 + ar) = 624$$

$$\Rightarrow ar \times 52 = 624 \quad \dots[\text{From (i)}]$$

$$\Rightarrow ar = 12$$

$$\Rightarrow a = \frac{12}{r}$$

Substituting in (i), we get

$$\frac{12}{r} + \frac{12}{r} \times r + \frac{12}{r} \times r^2 = 52$$

$$\Rightarrow \frac{12}{r} + 12 + 12r = 52$$

$$\Rightarrow \frac{12 + 12r + 12r^2}{r} = 52$$

$$\Rightarrow 12 + 12r + 12r^2 = 52r$$

$$\Rightarrow 12r^2 - 40r + 12 = 0$$

$$\Rightarrow 3r^2 - 10r + 3 = 0$$

$$\Rightarrow 3r^2 - 9r - r + 3 = 0$$

$$\Rightarrow 3r(r - 3) - 1(r - 3) = 0$$

$$\Rightarrow (3r - 1)(r - 3) = 0$$

$$\Rightarrow r = \frac{1}{3} \text{ or } r = 3$$

$$\Rightarrow a = \frac{12}{\frac{1}{3}} = 36 \text{ or } 4$$

Thus, required terms are:

$$\begin{aligned} a, ar, ar^2 &= 36, 36 \times \frac{1}{3}, 36 \times \frac{1}{9} \quad \text{OR} \quad 4, 4 \times 3, 4 \times 9 \\ &= 36, 12, 4 \quad \quad \quad \text{OR} \quad 4, 12, 36 \end{aligned}$$

### Question 8.

The sum of three numbers in G.P. is 21 and the sum of their squares is 189. Find the numbers.

### Solution:

Let the numbers be  $a$ ,  $ar$  and  $ar^2$ .

$$\Rightarrow (a)^2 + (ar)^2 + (ar^2)^2 = 189$$

$$\Rightarrow a^2 + a^2r^2 + a^2r^4 = 189$$

$$\text{And, } a + ar + ar^2 = 21$$

$$\Rightarrow (a + ar + ar^2)^2 = 21^2$$

$$\Rightarrow a^2 + a^2r^2 + a^2r^4 + 2a^2r + 2a^2r^3 + 2a^2r^2 = 441$$

$$\Rightarrow 189 + 2ar(a + ar^2 + ar) = 441$$

$$\Rightarrow 2ar \times 21 = 441 - 189$$

$$\Rightarrow 42ar = 252$$

$$\Rightarrow ar = 6$$

$$\Rightarrow r = \frac{6}{a}$$

$$\text{Now, } a + ar + ar^2 = 21$$

$$\Rightarrow a + a \times \frac{6}{a} + a \times \frac{36}{a^2} = 21$$

$$\Rightarrow a + 6 + \frac{36}{a} = 21$$

$$\Rightarrow a^2 + 6a + 36 = 21a$$

$$\Rightarrow a^2 - 15a + 36 = 0$$

$$\Rightarrow a^2 - 12a - 3a + 36 = 0$$

$$\Rightarrow a(a-12) - 3(a-12) = 0$$

$$\Rightarrow (a-12)(a-3) = 0$$

$$\Rightarrow a = 12 \text{ or } a = 3$$

$$\Rightarrow r = \frac{6}{12} = \frac{1}{2} \text{ or } r \frac{6}{3} = 2$$

Thus, required terms are:

$$a, ar, ar^2 = 12, 12 \times \frac{1}{2}, 12 \times \frac{1}{4} \quad \text{OR} \quad 3, 3 \times 2, 3 \times 4$$
$$= 12, 6, 3 \quad \text{OR} \quad 3, 6, 12$$